

TABLE OF CONTENTS

1. OVERVIEW	6
2. ECI REPORTING GUIDELINES	8
3. ECI CATEGORIES AND TERMS	8
3.1 Pneumonitis	9
3.2 Colitis	10
3.3 Endocrine	12
3.4 Hematologic	15
3.5 Hepatic	16
3.6 Neurologic	18
3.7 Ocular	19
3.8 Renal	20
3.9 Skin	21
3.9.1. Immediate Evaluation for Potential Skin ECIs	22
3.10 Other	23
3.11 Infusion Reactions	24
3.12 Follow-up to Resolution	26
4. REFERENCES	27
5. APPENDIX 1 –Events of Clinical Interest (ECI) – Reference Table	28
6. APPENDIX 2 – Past Medical History Related to Dermatologic Event	29
7. APPENDIX 3 – Presentation of the Dermatologic Event	30
8. APPENDIX 4 – Focused Skin Examination	32

1. OVERVIEW

The purpose of this document is to provide study sites with guidance on the identification and management of Events of Clinical Interest for the MK-3475 (also known as pembrolizumab) program.

Based on the literature review [1-11], and consideration of mechanism of action of pembrolizumab, potential immune-related adverse events (irAEs) are the primary Event of Clinical Interest (ECI). Immune-related AEs are adverse events associated with the treatment of patients with immunotherapy treatments that appear to be associated with the immune therapy's mechanism of action. Based on these potential irAEs, the sponsor has defined a list of specific adverse event terms (ECIs) that are selected adverse experiences that **must be reported to Merck within 24 hours** from the time the Investigator/physician is aware of such an occurrence, regardless of whether or not the investigator/physician considers the event to be related to study drug(s). In addition, these ECIs require additional detailed information to be collected and entered in the study database. ECIs may be identified through spontaneous patient report and / or upon review of subject data. **Table 1** provides the list of terms and reporting requirements for AEs that must be reported as ECIs for MK-3475 protocols. Of note, the requirement for reporting of ECIs applies to all arms, including comparators, of MK-3475 clinical trials

Given that our current list of events of clinical interest is not comprehensive for all potential immune-related events, it is possible that AEs other than those listed in this document may be observed in patients receiving pembrolizumab. Therefore any Grade 3 or higher event that the investigator/physician considers to be immune-related should be reported as an ECI regardless of whether the specific event term is in Table 1 and **reported to Merck within 24 hours** from the time the Investigator/physician is aware of such an occurrence. Adverse events that are both an SAE and an ECI should be reported one time as an SAE only, however the event must be appropriately identified as an ECI as well in the database.

Pembrolizumab Event of Clinical Interest Guidance Document

Table 1: Events of Clinical Interest

Pneumonitis (reported as ECI if ≥ Grade 2)		
Acute interstitial pneumonitis	Interstitial lung disease	Pneumonitis
Colitis (reported as ECI if > Grade 2 or any grade resulting in dose modification or use of systemic steroids to treat the AE)		
Intestinal Obstruction	Colitis	Colitis microscopic
Enterocolitis	Enterocolitis hemorrhagic	Gastrointestinal perforation
Necrotizing colitis	Diarrhea	
Endocrine (reported as ECI if ≥ Grade 3 or ≥ Grade 2 and resulting in dose modification or use of systemic steroids to treat the AE)		
Adrenal Insufficiency	Hyperthyroidism	Hypophysitis
Hypopituitarism	Hypothyroidism	Thyroid disorder
Thyroiditis	Hyperglycemia, if ≥ Grade 3 and associated with ketosis or metabolic acidosis (DKA)	
Endocrine (reported as ECI)		
Type 1 diabetes mellitus (if new onset)		
Hematologic (reported as ECI if ≥ Grade 3 or any grade resulting in dose modification or use of systemic steroids to treat the AE)		
Autoimmune hemolytic anemia	Aplastic anemia	Thrombotic Thrombocytopenic Purpura (TTP)
Idiopathic (or immune) Thrombocytopenia Purpura (ITP)	Disseminated Intravascular Coagulation (DIC)	Haemolytic Uraemic Syndrome (HUS)
Any Grade 4 anemia regardless of underlying mechanism		
Hepatic (reported as ECI if ≥ Grade 2, or any grade resulting in dose modification or use of systemic steroids to treat the AE)		
Hepatitis	Autoimmune hepatitis	Transaminase elevations (ALT and/or AST)
Infusion Reactions (reported as ECI for any grade)		
Allergic reaction	Anaphylaxis	Cytokine release syndrome
Serum sickness	Infusion reactions	Infusion-like reactions
Neurologic (reported as ECI for any grade)		
Autoimmune neuropathy	Guillain-Barre syndrome	Demyelinating polyneuropathy
Myasthenic syndrome		
Ocular (report as ECI if ≥ Grade 2 or any grade resulting in dose modification or use of systemic steroids to treat the AE)		
Uveitis	Iritis	
Renal (reported as ECI if ≥ Grade 2)		
Nephritis	Nephritis autoimmune	Renal Failure
Renal failure acute	Creatinine elevations (report as ECI if ≥ Grade 3 or any grade resulting in dose modification or use of systemic steroids to treat the AE)	
Skin (reported as ECI for any grade)		
Dermatitis exfoliative	Erythema multiforme	Stevens-Johnson syndrome
Toxic epidermal necrolysis		
Skin (reported as ECI if ≥ Grade 3)		
Pruritus	Rash	Rash generalized
Rash maculo-papular		
Any rash considered clinically significant in the physician's judgment		
Other (reported as ECI for any grade)		
Myocarditis	Pancreatitis	Pericarditis
Any other Grade 3 event which is considered immune-related by the physician		

Pembrolizumab Event of Clinical Interest Guidance Document

Each of the events above is described within this guidance document, along with site requirements for reporting these events to the Sponsor. The information collected should be entered into the narrative field(s) of the Adverse Event module in the database (please note, if narrative entry into the database is not available, please use the narrative text box on the 1727/AER Form). If additional Medical History or Concomitant Medications are reported, the Medical History and Concomitant Medication modules in the database must be updated.

In addition, the guidelines include recommendations on the management of these ECIs. These guidelines are intended to be applied when the physician determines the events to be related to pembrolizumab. **Note:** if after the evaluation the event is determined not to be related, the physician is instructed to follow the ECI reporting guidance but does not need to follow the treatment guidance (below). Therefore, these recommendations should be seen as guidelines and the treating physician should exercise individual clinical judgment based on the patient. For any question of dose modification or other treatment options, the specific language in the protocol should be followed. Any questions pertaining to the collection of this information or management of ECIs should be directed to your local Sponsor contact.

Dose Modification/Discontinuation

The treatment guidance provides specific direction when to hold and/or discontinue pembrolizumab for each immune related adverse event. Of note, when the guidance states to “discontinue” pembrolizumab this is the permanent discontinuation of treatment with pembrolizumab. “Hold” means to stop treating with pembrolizumab but resumption of treatment may be considered assuming the patient meets the criteria for resumption of treatment.

2. ECI REPORTING GUIDELINES

ECIs are selected non-serious and serious adverse experiences that must be reported to Merck **within 24 hours** regardless of attribution to study treatment. The AEs listed in this document and any event that meets the ECI criteria (as noted) in Table 1 or in the respective protocol (event term and Grade) **must be reported regardless of physician-determined causality with study medication and whether or not considered immune-related by the physician** (unless otherwise specified). Physicians/study coordinators/designated site personnel are required to record these experiences as ECIs on the Adverse Experience electronic Case Report Forms (eCRFs) (or on paper) and to provide supplemental information (such as medical history, concomitant medications, investigations, etc.) about the event.

- Please refer to the Data Entry Guidelines (DEGs) for your protocol.
- Please refer to protocol for details on reporting timelines and reporting of Overdose and Drug Induced Liver Injury (DILI).

3. ECI CATEGORIES AND TERMS

This section describes the ECI categories and outlines subject management guidelines when an ECI is reported.

3.1 Pneumonitis

The following AE terms, if considered \geq Grade 2, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Pneumonitis
- Interstitial lung disease
- Acute interstitial pneumonitis

If symptoms indicate possible new or worsening cardiac abnormalities additional testing and/or a cardiology consultation should be considered.

All attempts should be made to rule out other causes such as metastatic disease, bacterial or viral infection. It is important that patients with a suspected diagnosis of pneumonitis be managed as per the guidance below until treatment-related pneumonitis is excluded. Treatment of both a potential infectious etiology and pneumonitis in parallel may be warranted. Management of the treatment of suspected pneumonitis with steroid treatment should not be delayed for a therapeutic trial of antibiotics. If an alternative diagnosis is established, the patient does not require management as below; however the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Report as ECI
- Hold pembrolizumab.
- Consider pulmonary consultation with bronchoscopy and biopsy/BAL.
- Consider ID consult
- Conduct an in person evaluation approximately twice per week
- Consider frequent Chest X-ray as part of monitoring
- Treat with systemic corticosteroids at a dose of 1 to 2 mg/kg/day prednisone or equivalent. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.
- Second episode of pneumonitis – discontinue pembrolizumab if upon re-challenge the patient develops a second episode of Grade 2 or higher pneumonitis.

Grade 3 and 4 events:

- Report as ECI
- Discontinue pembrolizumab.
- Hospitalize patient
- Bronchoscopy with biopsy and/or BAL is recommended.
- Immediately treat with intravenous steroids (methylprednisolone 125 mg IV). When symptoms improve to Grade 1 or less, a high dose oral steroid (prednisone 1 to 2 mg/kg once per day or dexamethasone 4 mg every 4 hours) taper should be started and continued over no less than 4 weeks.
- If IV steroids followed by high dose oral steroids does not reduce initial symptoms within 48 to 72 hours, treat with additional anti-inflammatory measures. Discontinue additional anti-inflammatory measures upon symptom relief and initiate a prolonged steroid taper over 45 to 60 days. If symptoms worsen during steroid reduction, initiate a retapering of steroids starting at a higher dose of 80 or 100 mg followed by a more prolonged taper and administer additional anti-inflammatory measures, as needed
- Add prophylactic antibiotics for opportunistic infections.

3.2 Colitis

The following AE terms, if considered \geq Grade 2 or resulting in dose modification or use of systemic steroids to treat the AE, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Colitis
- Colitis microscopic
- Enterocolitis
- Enterocolitis hemorrhagic
- Gastrointestinal perforation
- Intestinal obstruction
- Necrotizing colitis
- Diarrhea

All attempts should be made to rule out other causes such as metastatic disease, bacterial or parasitic infection, viral gastroenteritis, or the first manifestation of an inflammatory bowel disease by examination for stool leukocytes, stool cultures, a *Clostridium difficile* titer and endoscopy. However the AE should be reported regardless of etiology.

Course of Action

Grade 2 Diarrhea/Colitis (4-6 stools/day over baseline, dehydration requiring IV fluids < 24 hours, abdominal pain, mucus or blood in stool):

- Report as ECI
- Hold pembrolizumab.
- Symptomatic Treatment
- For Grade 2 diarrhea that persists for greater than 3 days, and for diarrhea with blood and/or mucus,
 - o Consider GI consultation and endoscopy to confirm or rule out colitis
 - o Administer oral corticosteroids (prednisone 1-2 mg/kg QD or equivalent)
- When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.
- If symptoms worsen or persist > 3 days treat as Grade 3

Grade 3 Diarrhea/Colitis (or Grade 2 diarrhea that persists for > 1 week):

- Report as ECI
- Hold pembrolizumab.
- Rule out bowel perforation. Imaging with plain films or CT can be useful.
- Recommend consultation with Gastroenterologist and confirmation biopsy with endoscopy.
- Treat with intravenous steroids (methylprednisolone 125 mg) followed by high dose oral steroids (prednisone 1 to 2 mg/kg once per day or dexamethasone 4 mg every 4 hours) When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks. Taper over 6 to 8 weeks in patients with diffuse and severe ulceration and/or bleeding.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.
- If IV steroids followed by high dose oral steroids does not reduce initial symptoms within 48 to 72 hours, consider treatment with additional anti-inflammatory measures as described in the literature [5]. Discontinue additional anti-inflammatory measures upon symptom relief and initiate a prolonged steroid taper over 45 to 60 days. If symptoms worsen during steroid reduction, initiate a retapering of steroids starting at a higher dose of 80 or 100 mg followed by a more prolonged taper and administer additional anti-inflammatory measures as needed.

Pembrolizumab Event of Clinical Interest Guidance Document

Grade 4 events:

- Report as ECI
- Permanently discontinue pembrolizumab.
- Manage as per Grade 3.

3.3 Endocrine

The following AE terms, if considered \geq Grade 3 or if \geq Grade 2 and require holding/discontinuation/modification of pembrolizumab dosing, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Adrenal insufficiency
- Hyperthyroidism
- Hypophysitis
- Hypopituitarism
- Hypothyroidism
- Thyroid disorder
- Thyroiditis

All attempts should be made to rule out other causes such as brain metastases, sepsis and/or infection. However the AE should be reported regardless of etiology.

Hypophysitis or other symptomatic endocrinopathy other than hypo- or hyperthyroidism

Grade 2-4 events:

- Report as ECI if appropriate
- Hold pembrolizumab
- Rule out infection and sepsis with appropriate cultures and imaging.
- Monitor thyroid function or other hormonal level tests and serum chemistries more frequently until returned to baseline values.
- Pituitary gland imaging should be considered (MRIs with gadolinium and selective cuts of the pituitary can show enlargement or heterogeneity and confirm the diagnosis).
- Treat with prednisone 40 mg p.o. or equivalent per day. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks. Replacement of appropriate hormones may be required as the steroid dose is tapered.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.
- Hypophysitis with clinically significant adrenal insufficiency and hypotension, dehydration, and electrolyte abnormalities (such as hyponatremia and hyperkalemia) constitutes adrenal crisis.
- Consultation with an endocrinologist may be considered.

Pembrolizumab Event of Clinical Interest Guidance Document

Hyperthyroidism and Hypothyroidism

Thyroid disorders can occur at any time during treatment. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders.

Grade 2 hyperthyroidism, Grade 2-4 hypothyroidism events:

- Report as ECI if appropriate (see Table 1)
- Monitor thyroid function or other hormonal level tests and serum chemistries more frequently until returned to baseline values.
- Thyroid hormone and/or steroid replacement therapy to manage adrenal insufficiency.
- Therapy with pembrolizumab can be continued while treatment for the thyroid disorder is instituted.
- In hyperthyroidism, non-selective beta-blockers (e.g. propranolol) are suggested as initial therapy.
- In hypothyroidism, thyroid hormone replacement therapy, with levothyroxine or liothyronine, is indicated per standard of care.
- Consultation with an endocrinologist may be considered.

Grade 3 hyperthyroidism events:

- Report as ECI
- Hold pembrolizumab.
- Rule out infection and sepsis with appropriate cultures and imaging.
- Treat with an initial dose of methylprednisolone 1 to 2 mg/kg intravenously followed by oral prednisone 1 to 2 mg/kg per day. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks. Replacement of appropriate hormones may be required as the steroid dose is tapered.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 4 hyperthyroidism events:

- Report as ECI
- Discontinue pembrolizumab.
- Manage as per Grade 3

Pembrolizumab Event of Clinical Interest Guidance Document

Type 1 diabetes mellitus (if new onset) and \geq Grade 3 Hyperglycemia

The following AE terms are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Type 1 diabetes mellitus (T1DM), if new onset, including diabetic ketoacidosis (DKA)
- Grade 3 or higher hyperglycemia, if associated with ketosis (ketonuria) or metabolic acidosis (DKA).

Immune-mediated diabetes may present as new onset of Type 1 diabetes or an abrupt worsening of pre-existing diabetes associated with laboratorial evidence of beta cell failure. All attempts should be made to rule out other causes such as type 2 diabetes mellitus (T2DM), T2DM decompensation, steroid-induced diabetes, physiologic stress-induced diabetes, or poorly controlled pre-existing diabetes (either T1DM or T2DM), but events meeting the above criteria should be reported as ECIs regardless of etiology. The patients may present with hyperglycemia (abrupt onset or abrupt decompensation) with clinical evidence of diabetic ketoacidosis or laboratory evidence of insulin deficiency, such as ketonuria, laboratory evidence of metabolic acidosis, or low or undetected c-peptide.

Course of Action

T1DM should be immediately treated with insulin.

T1DM or Grade 3-4 Hyperglycemia events:

- Report as ECI if appropriate (see Table 1)
- Hold pembrolizumab for new onset Type 1 diabetes mellitus or Grade 3-4 hyperglycemia associated with evidence of beta cell failure, and resume pembrolizumab when patients are clinically and metabolically stable.
- Insulin replacement therapy is recommended for Type 1 diabetes mellitus and for Grade 3-4 hyperglycemia associated with metabolic acidosis or ketonuria.
- Evaluate patients with serum glucose and a metabolic panel, urine ketones, glycosylated hemoglobin, and C-peptide.
- Consultation with an Endocrinologist is recommended.
- Consider local testing for islet cell antibodies and antibodies to GAD, IA-2, ZnT8, and insulin may be obtained.

Pembrolizumab Event of Clinical Interest Guidance Document

3.4 Hematologic

The following AE term, if considered Grade ≥ 3 or requiring dose modification or use of systemic steroids to treat the AE, are considered an ECI and should be reported to the Sponsor within 24 hours of the event:

- Autoimmune hemolytic anemia
- Aplastic anemia
- Disseminated Intravascular Coagulation (DIC)
- Haemolytic Uraemic Syndrome (HUS)
- Idiopathic (or immune) Thrombocytopenia Purpura (ITP)
- Thrombotic Thrombocytopenic Purpura (TTP)
- Any Grade 4 anemia regardless of underlying mechanism

All attempts should be made to rule out other causes such as metastases, sepsis and/or infection. Relevant diagnostic studies such as peripheral blood smear, reticulocyte count, LDH, haptoglobin, bone marrow biopsy or Coomb's test, etc., should be considered to confirm the diagnosis. However the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Report as ECI
- Hold pembrolizumab
- Prednisone 1-2 mg/kg daily may be indicated
- Consider Hematology consultation.
Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 3 events:

- Report as ECI
- Hematology consultation.
- Hold pembrolizumab Discontinuation should be considered as per specific protocol guidance.
- Treat with methylprednisolone 125 mg iv or prednisone 1-2 mg/kg p.o. (or equivalent) as appropriate
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 4 events:

- Report as ECI
- Hematology consultation
- Discontinue pembrolizumab for all solid tumor indications; refer to protocol for hematologic malignancies.
- Treat with methylprednisolone 125 mg iv or prednisone 1-2 mg/kg p.o. (or equivalent) as appropriate

3.5 Hepatic

The following AE terms, if considered \geq Grade 2 or greater (or any grade with dose modification or use of systemic steroids to treat the AE), are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Autoimmune hepatitis
- Hepatitis
- Transaminase elevations

All attempts should be made to rule out other causes such as metastatic disease, infection or other hepatic diseases. However the AE should be reported regardless of etiology.

Drug Induced Liver Injury (DILI)

In addition, the event must be reported as a Drug Induced Liver Injury (DILI) ECI, if the patient meets the laboratory criteria for potential DILI defined as:

- An elevated alanine transaminase (ALT) or aspartate transaminase (AST) lab value that is greater than or equal to three times (3X) the upper limit of normal (ULN) and
- An elevated total bilirubin lab value that is greater than or equal to two times (2X) ULN and
- At the same time, an alkaline phosphatase (ALP) lab value that is less than 2X ULN,
- As a result of within-protocol-specific testing or unscheduled testing.

Note that any hepatic immune ECI meeting DILI criteria should only be reported once as a DILI event.

Course of Action

Grade 2 events:

- Report as ECI
- Hold pembrolizumab when AST or ALT >3.0 to 5.0 times ULN and/or total bilirubin >1.5 to 3.0 times ULN.
- Monitor liver function tests more frequently until returned to baseline values (consider weekly).
 - o Treat with 0.5 - 1 mg/kg/day methylprednisolone or oral equivalent and when LFT returns to grade 1 or baseline, taper steroids over at least 1 month, consider prophylactic antibiotics for opportunistic infections, and resume pembrolizumab per protocol
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.
- Permanently discontinue pembrolizumab for patients with liver metastasis who begin treatment with Grade 2 elevation of AST or ALT, and AST or ALT increases $\geq 50\%$ relative to baseline and lasts ≥ 1 week.

Grade 3 events:

- Report as ECI
- Discontinue pembrolizumab when AST or ALT >5.0 times ULN and/or total bilirubin >3.0 times ULN.
- Consider appropriate consultation and liver biopsy to establish etiology of hepatic injury, if necessary
- Treat with high-dose intravenous glucocorticosteroids for 24 to 48 hours. When symptoms improve to Grade 1 or less, a steroid taper with dexamethasone 4 mg every 4 hours or prednisone at 1 to 2 mg/kg should be started and continued over no less than 4 weeks.
- If serum transaminase levels do not decrease 48 hours after initiation of systemic steroids, oral mycophenolate mofetil 500 mg every 12 hours may be given. Infliximab is not recommended due to its potential for hepatotoxicity.

Pembrolizumab Event of Clinical Interest Guidance Document

- Several courses of steroid tapering may be necessary as symptoms may worsen when the steroid dose is decreased.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 4 events:

- Report as ECI
- Permanently discontinue pembrolizumab
- Manage patient as per Grade 3 above

Pembrolizumab Event of Clinical Interest Guidance Document

3.6 Neurologic

The following AE terms, regardless of grade, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Autoimmune neuropathy
- Demyelinating polyneuropathy
- Guillain-Barre syndrome
- Myasthenic syndrome

All attempts should be made to rule out other causes such as metastatic disease, other medications or infectious causes. However the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Report as ECI
- Moderate (Grade 2) – consider withholding pembrolizumab.
- Consider treatment with prednisone 1-2 mg/kg p.o. daily as appropriate
- Consider Neurology consultation. Consider biopsy for confirmation of diagnosis.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 3 and 4 events:

- Report as ECI
- Discontinue pembrolizumab
- Obtain neurology consultation. Consider biopsy for confirmation of diagnosis
- Treat with systemic corticosteroids at a dose of 1 to 2 mg/kg prednisone or equivalent once per day. If condition worsens consider IVIG or other immunosuppressive therapies as per local guidelines

When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.

Pembrolizumab Event of Clinical Interest Guidance Document

3.7 Ocular

The following AE terms, if considered Grade ≥ 2 or requiring dose modification or use of systemic steroids to treat the AE, is considered an ECI and should be reported to the Sponsor within 24 hours of the event:

- Uveitis
- Iritis

All attempts should be made to rule out other causes such as metastatic disease, infection or other ocular disease (e.g. glaucoma or cataracts). However the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Evaluation by an ophthalmologist is strongly recommended.
- Treat with topical steroids such as 1% prednisolone acetate suspension and iridocyclitics.
- Discontinue pembrolizumab as per protocol if symptoms persist despite treatment with topical immunosuppressive therapy.

Grade 3 events:

- Evaluation by an ophthalmologist is strongly recommended
- Hold pembrolizumab and consider permanent discontinuation as per specific protocol guidance.
- Treat with systemic corticosteroids such as prednisone at a dose of 1 to 2 mg/kg per day. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 4 events:

- Evaluation by an ophthalmologist is strongly recommended
- Permanently discontinue pembrolizumab.
- Treat with corticosteroids as per Grade 3 above

Pembrolizumab Event of Clinical Interest Guidance Document

3.8 Renal

The following AEs if \geq Grade 2 are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Nephritis
- Nephritis autoimmune
- Renal failure
- Renal failure acute

Creatinine elevations \geq Grade 3 or any grade with dose modification or use of systemic steroids to treat the AE.

All attempts should be made to rule out other causes such as obstructive uropathy, progression of disease, or injury due to other chemotherapy agents. A renal consultation is recommended. However the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Hold pembrolizumab
- Treatment with prednisone 1-2 mg/kg p.o. daily.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 3-4 events:

- Discontinue pembrolizumab
- Renal consultation with consideration of ultrasound and/or biopsy as appropriate
- Treat with systemic corticosteroids at a dose of 1 to 2 mg/kg prednisone IV or equivalent once per day.

When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.

Pembrolizumab Event of Clinical Interest Guidance Document

3.9 Skin

Rash and Pruritus

The following AEs should be considered as ECIs, if \geq Grade 3 and should be reported to the Sponsor within 24 hours of the event:

- Pruritus
- Rash
- Rash generalized
- Rash maculo-papular
- In addition to CTCAE Grade 3 rash, any rash that is considered clinically significant, in the physician's judgment, should be treated as an ECI. Clinical significance is left to the physician to determine, and could possibly include rashes such as the following:
 - o rash with a duration >2 weeks; OR
 - o rash that is $>10\%$ body surface area; OR
 - o rash that causes significant discomfort not relieved by topical medication or temporary cessation of study drug.

Other Skin ECIs

The following AEs should **always** be reported as ECIs, regardless of grade, and should be reported to the Sponsor within 24 hours of the event:

- Dermatitis exfoliative
- Erythema multiforme
- Steven's Johnson syndrome
- Toxic epidermal necrolysis

Please note, the AE should be reported regardless of etiology.

Course of Action

Grade 2 events:

- Symptomatic treatment should be given such as topical glucocorticosteroids (e.g., betamethasone 0.1% cream or hydrocortisone 1%) or urea-containing creams in combination with oral anti-pruritics (e.g., diphenhydramine HCl or hydroxyzine HCl).
- Treatment with oral steroids is at physician's discretion for Grade 2 events.

Grade 3 events:

- Hold pembrolizumab.
- Consider Dermatology Consultation and biopsy for confirmation of diagnosis.
- Treatment with oral steroids is recommended, starting with 1 mg/kg prednisone or equivalent once per day or dexamethasone 4 mg four times orally daily. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks.

Grade 4 events:

- Permanently discontinue pembrolizumab.
- Dermatology consultation and consideration of biopsy and clinical dermatology photograph.
- Initiate steroids at 1 to 2 mg/kg prednisone or equivalent. When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.

Pembrolizumab Event of Clinical Interest Guidance Document

3.9.1. Immediate Evaluation for Potential Skin ECIs

A. Photographs:

Every attempt should be made to get a photograph of the actual ECI skin lesion or rash as soon as possible. Obtain appropriate consent for subject photographs if a consent form addendum is required by your IRB/ERC.

- Take digital photographs of:
 - o the head (to assess mucosal or eye involvement),
 - o the trunk and extremities, and
 - o a close-up of the skin lesion/rash.
- If possible, a ruler should be placed alongside the site of a skin occurrence as a fixed marker of distance.
- The time/date stamp should be set in the 'ON' position for documentation purposes.
- Photographs should be stored with the subject's study records.
- The Sponsor may request copies of photographs. The local study contact (e.g., CRA) will provide guidance to the site, if needed.

B. Past Medical History:

Collect past medical history relevant to the event, using the questions in Appendix 2 (Past Medical History Related to Dermatologic Event) as a guide. Any preexisting conditions not previously reported (e.g., drug allergy) should be entered into the Medical History eCRF.

C. Presentation of the Event:

Collect information on clinical presentation and potential contributing factors using the questions in Appendix 3 (Presentation of the Dermatologic Event) as a guide. This information should be summarized and entered in narrative format in the AE eCRF. Please use the available free-text fields, such as Signs and Symptoms. Note pertinent negatives where applicable to reflect that the information was collected. Any treatments administered should be entered on the Concomitant Medication eCRF.

D. Vitals Signs and Standard Laboratory Tests:

Measure vital signs (pulse, sitting BP, oral temperature, and respiratory rate) and record on the Vital Signs eCRF. Perform standard laboratory tests (CBC with manual differential and serum chemistry panel, including LFTs).

E. Focused Skin Examination:

Perform a focused skin examination using the questions in Appendix 4 (Focused Skin Examination) as a guide. Information should be summarized and entered on the Adverse Experience eCRF as part of the narrative.

F. Dermatology Consult

Refer the subject to a dermatologist as soon as possible.

- For a "severe rash", the subject must be seen within 1-2 days of reporting the event.
- For clinically significant rash, the subject should be seen within 3-5 days.

The dermatologist should submit a biopsy sample to a certified dermatopathology laboratory or to a pathologist experienced in reviewing skin specimens.

The site should provide the dermatologist with all relevant case history, including copies of clinical photographs and laboratory test results.

3.10 Other

The following AEs, regardless of grade, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Myocarditis
- Pericarditis
- Pancreatitis
- Any additional Grade 3 or higher event which the physician considers to be immune related

All attempts should be made to rule out other causes. Therapeutic specialists should be consulted as appropriate. However the AE should be reported regardless of etiology.

Course of Action

Grade 2 events or Grade 1 events that do not improve with symptomatic treatment:

- Withhold pembrolizumab.
- Systemic corticosteroids may be indicated.
- Consider biopsy for confirmation of diagnosis.
- If pembrolizumab held and corticosteroid required, manage as per grade 3 below.

Grade 3 events:

- Hold pembrolizumab
- Treat with systemic corticosteroids at a dose of 1 to 2 mg/kg prednisone or equivalent once per day.
- When symptoms improve to Grade 1 or less, steroid taper should be started and continued over no less than 4 weeks.
- Permanently discontinue for inability to reduce corticosteroid dose to 10 mg or less of prednisone or equivalent per day within 12 weeks. Otherwise, pembrolizumab treatment may be restarted and the dose modified as specified in the protocol

Grade 4 events:

- Treat with systemic corticosteroids at a dose of 1 to 2 mg/kg prednisone or equivalent once per day.
- Discontinue pembrolizumab

3.11 Infusion Reactions

The following AE terms, regardless of grade, are considered ECIs and should be reported to the Sponsor within 24 hours of the event:

- Allergic reaction
- Anaphylaxis
- Cytokine release syndrome
- Serum sickness
- Infusion reactions
- Infusion-like reactions

Please note, the AE should be reported regardless of etiology.

Course of Action

Refer to infusion reaction table in the protocol and below.

Pembrolizumab Event of Clinical Interest Guidance Document

Infusion Reactions

NCI CTCAE Grade	Treatment	Premedication at subsequent dosing
<p>Grade 1 Mild reaction; infusion interruption not indicated; intervention not indicated</p>	<p>Increase monitoring of vital signs as medically indicated until the subject is deemed medically stable in the opinion of the investigator.</p>	<p>None</p>
<p>Grade 2 Requires infusion interruption but responds promptly to symptomatic treatment (e.g., antihistamines, NSAIDS, narcotics, IV fluids); prophylactic medications indicated for <=24 hrs</p>	<p>Stop Infusion. Additional appropriate medical therapy may include but is not limited to: IV fluids Antihistamines NSAIDS Acetaminophen Narcotics Increase monitoring of vital signs as medically indicated until the subject is deemed medically stable in the opinion of the investigator. If symptoms resolve within one hour of stopping drug infusion, the infusion may be restarted at 50% of the original infusion rate (e.g. from 100 mL/hr to 50 mL/hr). Otherwise dosing will be held until symptoms resolve and the subject should be premedicated for the next scheduled dose. Subjects who develop Grade 2 toxicity despite adequate premedication should be permanently discontinued from further trial treatment administration.</p>	<p>Subject may be premedicated 1.5h (± 30 minutes) prior to infusion of pembrolizumab with: Diphenhydramine 50 mg p.o. (or equivalent dose of antihistamine). Acetaminophen 500-1000 mg p.o. (or equivalent dose of antipyretic).</p>
<p>Grades 3 or 4 Grade 3: Prolonged (i.e., not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalization indicated for other clinical sequelae (e.g., renal impairment, pulmonary infiltrates) Grade 4: Life-threatening; pressor or ventilatory support indicated</p>	<p>Stop Infusion. Additional appropriate medical therapy may include but is not limited to: IV fluids Antihistamines NSAIDS Acetaminophen Narcotics Oxygen Pressors Corticosteroids Epinephrine Increase monitoring of vital signs as medically indicated until the subject is deemed medically stable in the opinion of the investigator. Hospitalization may be indicated. Subject is permanently discontinued from further trial treatment administration.</p>	<p>No subsequent dosing</p>
<p>Appropriate resuscitation equipment should be available in the room and a physician readily available during the period of drug administration. For Further information, please refer to the Common Terminology Criteria for Adverse Events v4.0 (CTCAE) at http://ctep.cancer.gov</p>		

3.12 Follow-up to Resolution

Subjects should be followed to resolution. The Adverse Experience eCRF should be updated with information regarding duration and clinical course of the event. Information obtained from the consulting specialist, including diagnosis, should be recorded in the appropriate AE fields. Free-text fields should be used to record narrative information:

- Clinical course of the event
- Course of treatment
- Evidence supporting recovery
- Follow-up to the clinical course

Any treatments administered for the event should also be entered in the Concomitant Medication eCRF.

4. REFERENCES

1. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nature Reviews Cancer* 2012; 12:252-264.
2. Weber JS Practical Management of Immune-Related Adverse Events from Immune Checkpoint Protein Antibodies for the Oncologist. *American Society of Clinical Oncology* 2012; 1092-9118/10/1-10.
3. Weber JS, Kaehler KC, and Hauschild A. Management of Immune-Related Adverse Events and Kinetics of Response With Ipilimumab. *J Clin Oncol* 30. 2012. <http://jco.ascopubs.org/cgi/doi/10.1200/JCO.2012.41.6750>.
4. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, et al. Improved Survival with Ipilimumab in Patients with Metastatic Melanoma. *N Engl J Med* 363:711-723, 2010.
5. Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, et al. Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer. *N Engl J Med* 2012; 366:2443-2454.
6. Brahmer JR, Tykodi SS, Chow LQM, Hwu WJ, et al. Safety and Activity of Anti-PD-L1 Antibody in Patients with Advanced Cancer. *N Engl J Med* 2012; 366:2455-2465.
7. Weber J, Thompson JA, Hamid O, et al: A randomized, double-blind, placebo-controlled, phase II study comparing the tolerability and efficacy of ipilimumab administered with or without prophylactic budesonide in patients with unresectable stage III or IV melanoma. *Clin Cancer Res* 15:5591-5598, 2009.
8. Lemech C and Arkenau HT. Novel Treatments for Metastatic Cutaneous Melanoma and the Management of Emergent Toxicities. *Clinical Medicine Insights: Oncology* 2012;6 53-66.
9. Phan GQ, Weber JS, and Sondak VK. CTLA-4 Blockade with Monoclonal Antibodies in Patients with Metastatic Cancer: Surgical Issues. *Annals of Surgical Oncology* 15(11):3014-3021.
10. Bristol-Myers Squibb: YERVOY (ipilimumab): Serious and fatal immune-mediated adverse reactions—YERVOY Risk Evaluation and Mitigation Strategy (REMS). <http://www.yervoy.com/hcp/rem.s.aspx>
11. Bristol-Myers Squibb: YERVOY (ipilimumab) prescribing information revised March 2011. http://www.accessdata.fda.gov/drugsatfda_docs/label/2011/125377s0000lbl.pdf